

長庚大學101學年度研究所碩士班(含在職專班)招生考試試題

系所：護理學系碩士班

考試科目：社區衛生護理學

注意：請詳細閱讀下列試題，並請標明題號依試題順序將答案書寫於答案卷上。 本試題共 2 頁：第 | 頁

1. 請描述健康城市的特性(10%)及推動健康城市之步驟 (15%)。

2. 請描述社區健康評估資料收集的方式與步驟(10%)和社區健康評估的項目 (15%)。

長庚大學101學年度研究所碩士班(含在職專班)招生考試試題

系所：護理學系碩士班

考試科目：社區衛生護理學

注意：請詳細閱讀下列試題，並請標明題號依試題順序將答案書寫於答案卷上。 本試題共~~二~~頁：第~~二~~頁

3. 請解釋何謂自我效能 (Self Efficacy) 及其機轉 (25%)。

4. 請描述及解釋下列表的意義 (25%)。

(摘自 Ng, T., Niti, M., Zaw, M. and Kua, E. (2009). Depressive Symptoms and Incident Cognitive Impairment in Cognitively Well-Functioning Older Men and Women. *J Am Geriatr Soc*, 57,1058–1063.)

Table 2. Association Between Depressive Symptoms and Incident Cognitive Impairment at Follow-Up in Men and Women Stratified According to Hypertension and Vascular Factor Status

| Hypertension and Vascular Factors | Men | | | Women | | | Interaction P-Value | |
|--------------------------------------|--|------------------|---------|--|------------------|--------------|------------------------|--|
| | Subjects with Incident Cognitive Impairment/ Subjects in Sample n/N (%) | AOR (95% CI) | P-Value | Subjects with Incident Cognitive Impairment/ Subjects in Sample n/N (%) | AOR (95% CI) | P-Value | | |
| | | | | Overall | No hypertension | Hypertension | | |
| Not depressed | 9/485 (1.9) | 1.00 | | 26/844 (3.1) | 1.00 | | | |
| Depressed | 5/58 (8.6) | 4.74 (1.25–17.8) | .02 | 4/100 (4.0) | 1.29 (0.41–4.03) | .66 | .02 | |
| No hypertension | | | | | | | | |
| Not depressed | 4/208 (1.9) | 1.00 | | 14/397 (3.5) | 1.00 | | | |
| Depressed | 0/26 (0) | NA | | 0/50 (0) | NA | | | |
| Hypertension | | | | | | | | |
| Not depressed | 5/277 (1.8) | 1.00 | | 12/447 (2.7) | 1.00 | | | |
| Depressed | 5/32 (15.6) | 5.07 (1.06–24.2) | .04 | 4/50 (8.0) | 2.88 (0.82–10.2) | .10 | .02* | |
| No vascular factors | | | | | | | | |
| Not depressed | 3/97 (3.1) | 1.00 | | 4/184 (2.2) | 1.00 | | | |
| Depressed | 0/10 (0.0) | NA | | 0/17 (0.0) | NA | | | |
| Vascular factors | | | | | | | | |
| Not depressed | 6/388 (1.5) | 1.00 | | 22/660 (3.3) | 1.00 | | | |
| Depressed | 5/48 (10.4) | 4.77 (1.08–21.1) | .04 | 4/83 (4.8) | 1.62 (0.51–5.15) | .42 | .01* | |

Subjects who were not cognitively impaired (Mini-Mental State Examination (MMSE) score ≥ 24) at baseline were followed up and reassessed after 1.5 to 2 years. Cognitive impairment at follow-up was defined as MMSE score < 24 .

Adjusted for age, education level, apolipoprotein E e4 status, length of follow-up, and baseline MMSE score.

* Tests whether the effect of depressive symptoms on cognitive impairment incidence differs by sex in each stratum of hypertension status or vascular factor stratum.
AOR = adjusted odds ratio; CI = confidence interval; NA = not applicable.