

# 國立中山大學 112 學年度 碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

## — 作答注意事項 —

考試時間：100 分鐘

- 考試開始鈴響前不得翻閱試題，並不得書寫、劃記、作答。請先檢查答案卷（卡）之應考證號碼、桌角號碼、應試科目是否正確，如有不同立即請監試人員處理。
- 答案卷限用藍、黑色筆(含鉛筆)書寫、繪圖或標示，可攜帶橡皮擦、無色透明無文字墊板、尺規、修正液（帶）、手錶(未附計算器者)。每人每節限使用一份答案卷，請衡酌作答(不得另攜帶紙張，亦不得使用應考證空白處作為計算紙使用)。
- 答案卡請以 2B 鉛筆劃記，不可使用修正液（帶）塗改，未使用 2B 鉛筆、劃記太輕或污損致光學閱讀機無法辨識答案者，後果由考生自負。
- 答案卷（卡）應保持清潔完整，不得折疊、破壞或塗改應考證號碼及條碼，亦不得書寫考生姓名、應考證號碼或與答案無關之任何文字或符號。
- 可否使用計算機請依試題資訊內標註為準，如「可以」使用，廠牌、功能不拘，唯不得攜帶具有通訊、記憶或收發等功能或其他有礙試場安寧、考試公平之各類器材、物品（如鬧鈴、行動電話、電子字典等）入場。
- 試題及答案卷（卡）請務必繳回，未繳回者該科成績以零分計算。
- 試題採雙面列印，考生應注意試題頁數確實作答。
- 違規者依本校招生考試試場規則及違規處理辦法處理。

國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 1 頁

物理化學

Useful numbers:

molar gas constant  $R = 8.314 \text{ J K}^{-1} \text{ mole}^{-1} = 0.08314 \text{ L bar K}^{-1} \text{ mole}^{-1} = 0.08206 \text{ L atm K}^{-1} \text{ mole}^{-1}$

Boltzmann's constant  $k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$

Avogadro's number  $N_A = 6.02 \times 10^{23} \text{ mole}^{-1}$

Planck constant  $h = 6.63 \times 10^{-34} \text{ J sec}$

1 atm = 760 torr =  $1.013 \times 10^5 \text{ Pa}$ , 1 bar = 750 torr =  $10^5 \text{ Pa}$  ( $\text{Nt m}^{-2}$ )

Speed of light  $c = 299792458 \text{ m sec}^{-1}$

單選題：(每題 2 分，共 50 分)

(1) Given a free particle in a box, which of the following quantum mechanical conditions is not true?

(A) Shifting the box in space will change the wavefunction.

(B) The precise position of the particle is unpredictable.

(C) If the size of the box is infinite, the energy of the particle is not quantized.

(D) Shifting the box in space will change the energy of the particle.

(E) The energy of the particle depends on the shape of the box.

(2) Which one of the following is the correct form for the lowest-energy eigenfunction for a particle in a 1D box with infinite barriers at  $x = -L/2$  and  $L/2$ ?


(A)  $\sqrt{\frac{2}{L}} \sin(\pi x / L)$  (B)  $\sqrt{\frac{2}{L}} \cos(\pi x / L)$  (C)  $\sqrt{\frac{2}{L}} \frac{1}{i} (e^{i\pi x / L} - e^{-i\pi x / L})$  (D)  $\sqrt{\frac{2}{L}} (e^{i\pi x / L} + e^{-i\pi x / L})$  (E)

None of the above.

(3) Given a wave function  $\psi(x) = Nx(x-1)$  in the region  $0 \leq x \leq 1$ . Where  $N$  is the normalization constant to be determined. What is the possible value of  $N$ ?

(A)  $\frac{1}{\sqrt{30}}$  (B)  $\sqrt{2}$  (C)  $\sqrt{1/2}$  (D)  $\sqrt{30}$  (E) None of the above.

(4) The wavefunction on the right illustrates the quantum state of a harmonic oscillator with its vibrational quantum number  $\nu =$  (A) 1 (B) 2 (C) 3 (D) 4 (E) 5.



國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 2 頁

(5) Which of the following statements is true for a given set of harmonic oscillator wavefunctions corresponding to the same potential  $V(x)$  ?

- (A) The ground-state energy is called zero-point energy and its value is zero.
- (B) The mean displacement from the equilibrium distance  $\langle x \rangle \neq 0$  for the first excited vibrational state  $v = 1$ .
- (C) The wavefunctions must have zero amplitude beyond the classical turning points.
- (D)  $\int_{-\infty}^{\infty} \psi_v^* \psi_{v'} dx = 0, v \neq v'$
- (E) The wavefunctions involve Gaussian-type functions and thus are all even functions.

(6) Which of the following about the spherical harmonics  $Y_l^m(\theta, \phi)$  is not true?

- (A) The number of angular nodes in  $Y_l^m$  is  $l$ .
- (B) The complex  $Y_l^m$  give non-zero eigenvalues of  $\hat{L}_z$  which represents the operator of the  $z$ -component of angular momentum.
- (C)  $Y_0^0$  is a constant everywhere in space.
- (D)  $Y_l^m$  serves as eigenfunctions for  $\hat{L}_x$ .
- (E)  $Y_1^0 = \left(\frac{3}{4\pi}\right)^{1/2} \cos\theta$  correlates with all  $p_z$  orbitals.

(7) Which statement about H-atom is not true?

- (A) Spherical polar coordinates are the natural choice for H-atom Schrödinger equation because the potential term is spherically symmetric.
- (B) H-atom Schrödinger equation can be solved by separation of variables where the radial part  $R_{nl}(r)$  and the angular part  $Y_l^m(\theta, \phi) = \Theta(\theta) \Phi(\phi)$  are linked through  $l(l+1)$  while  $\Theta(\theta)$  and  $\Phi(\phi)$  are linked through  $m^2$ .
- (C) The operator for the  $z$ -component of angular momentum has the simplest form in spherical polar coordinates as opposed to the  $x$ - and  $y$ -components.
- (D) The rules for allowed angular momentum magnitudes and orientations are shared by the rigid rotor and the H-atom.
- (E) By changing the variable  $\sin\theta$  into  $x$ , the  $\Theta$  differential equation can be recognized as being the associated Legendre equation.


國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 3 頁



(8) In the above plot, the letters **a**, **b**, and **c** label three different radial distribution functions (RDF's)  $r^2R^2$  of H-atom from the same  $n$  level. Which of the following statements is true?

- (A) The maximum of curve **a** corresponds to the radii of maximum probability  $r_{mp} = n^2 a_0$  ( $a_0$  is Bohr radius).
- (B) Curve **b** corresponds to the RDF derived from a radial function  $R$  that is proportional to  $r^2 \exp(-r/3a_0)$ .
- (C) Curve **c** corresponds to angular momentum quantum number  $l = 2$ .
- (D) Curve **c** shows two angular nodes.
- (E) The principal quantum number  $n = 2$ .

(9) The term “intersystem crossing” is used for

- (A) a transition from an excited singlet state to another excited singlet state.
- (B) a transition from the ground singlet state to an excited singlet state.
- (C) a transition from an excited singlet state to an excited triplet state.
- (D) a transition from an excited triplet state to another excited triplet state.
- (E) a transition from an excited triplet state to the ground singlet state.

國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題


科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 4 頁

(10) Which of the following statements is not true for the experimental HCl rotational-vibrational spectrum below?



- (A) The splitting in each line is due the isotope effect.  
 (B) In the middle of the spectrum a forbidden line is missing by the selection rules.  
 (C) The bond length for HCl can be determined from the intervals between neighboring lines.  
 (D) The separation between lines decreases in the R-branch and increases in the P-branch.  
 (E) The lines connected to transitions with  $\Delta J = +1$  form the P-branch.
- (11) For which of the following thermodynamic quantities can an absolute value be calculated?  
 (A) internal energy  $U$  (B) enthalpy  $H$  (C) entropy  $S$  (D) Helmholtz energy  $A$  (E) Gibbs energy  $G$
- (12) For which of the following substances the standard molar entropy  $\bar{S}^\circ$  is zero at  $25^\circ\text{C}$ ?  
 (A)  $\text{Na}_{(s)}$  (B)  $\text{N}_{2(g)}$  (C)  $\text{I}_{2(s)}$  (D) all of these (E) None of these.
- (13) For which of the following standard enthalpies of formation is not zero  $\Delta\bar{H}_f^\circ(25^\circ\text{C})$ ?  
 (A)  $\text{Na}_{(s)}$  (B)  $\text{N}_{2(g)}$  (C)  $\text{I}_{2(g)}$  (D)  $\text{C}_{(s, \text{graphite})}$  (E) None of these.
- (14) For a reversible adiabatic expansion of an ideal gas, which of the following is always true?  
 (A) The work done by the gas is equal to the decrease in the internal energy of the gas.  
 (B) The work done by the gas is equal to the heat absorbed by the gas.  
 (C) The temperature of the gas will rise.  
 (D)  $PV=C$  where  $C$  is a constant along the path.  
 (E) No work is done by the gas to the surroundings.

國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 5 頁

(15) For which statement about the Joule-Thomson (J-T) experiment is not true?

(A) J-T expansion is an isenthalpic process ( $\Delta H = 0$ ).

(B)  $\left(\frac{\partial T}{\partial P}\right)_H$  is called the J-T coefficient  $\mu_{JT}$ .

(C) Any gas will be cooled by J-T expansion.

(D) For ideal gases  $\mu_{JT} = 0$ .

(E) A positive J-T coefficient implies that the gas cools upon expansion.

(16) For a gas whose equation of state is  $P(\bar{V} - b) = RT$  (a simplified *Van der Waals* equation), the generalized  $C_p - C_v = \frac{TV\alpha^2}{\kappa}$  ( $\alpha$  and  $\kappa$  represent thermal expansion coefficient and isothermal compressibility, respectively). The difference of these two heat capacities is equal to

(A)  $nR$  (B)  $nR\left(1 + \frac{\bar{V}}{b}\right)$  (C)  $R - b$  (D) 0 (E)  $nb$

(17) Which of the following is true?

(A) An increase of pressure always leads to a boiling point elevation for all substances.

(B) An increase of pressure always leads to a melting point depression for all substances.

(C) The slope of the solid-liquid coexistence curve in the P-T phase diagram is positive if the solid is less dense than the liquid.

(D) In the P-V phase diagram, the solid-liquid coexistence region is broader than the gas-liquid coexistence region.

(E) In the P-T phase diagram the Clausius-Clapeyron equation can be used to show that the slope of the solid-gas coexistence curve must be less than the slope of the liquid-gas coexistence curve near the triple point.

(18) Given that the molar enthalpy of vaporization is 42.7 kJ/mole and is independent of temperature, what is the boiling point of water if the atmospheric pressure becomes 427 torr?

(A) 85 °C (B) 77 °C (C) 56 °C (D) 42 °C (E) None of these.

(19) Which of the following about thermodynamic equilibria is not true?

(A) At phase equilibrium, a substance that occurs in two phases will have the same concentration in both phases.

國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 6 頁

(B) For a single component substance at a fixed value of pressure, the change of slopes in the plot of the chemical potential versus temperature across different phases represents the variation of the molar entropies of phases.

(C)  $\Delta_r G$  is the net difference of chemical potentials between products and reactants for the reaction.

(D)  $\Delta_r G^\circ$  representing the change in standard Gibbs energy for the reaction between unmixed reactants to form unmixed products is a function of temperature only.

(E)  $\Delta_r G = -RT \ln K_p + RT \ln Q$  where  $Q$  represents the reaction quotient.

(20) Which of the following statements is not true?

(A) The total partition function  $Q$  for a collection of  $N$  *independent* molecules or atoms can be expressed as the product of the molecular partition functions  $q$  for the individual species.

(B) If these molecules or atoms are *indistinguishable*, the total partition function  $Q$  is divided by a factor of  $N!$ .

(C) The existence of the  $1/N!$  factor in the expression of  $Q$  for ideal gases is noticeable only in the entropy expression, but is not noticeable in the internal energy expression because the differential of the logarithm of  $1/N!$  is zero.

(D) The molecular partition function  $q$  for an ideal gas molecule can be decomposed into the sum of the partition functions for four types of degree-of-freedom: translational, vibrational, rotational, and electronic.

(E) The vibrational partition function of a polyatomic molecule is a product of harmonic oscillator partition functions for each normal mode.

(21) Which of the following statements is not true?

(A) The most probable component of speed  $u_x$  of a gas is zero.

(B) The  $\langle u_x \rangle$  of a gas is zero.

(C) The broadness of the distribution function  $f(u_x)$  correlates with  $1/\sqrt{T}$ .

(D) If two different gases are at the same temperature, the probability distribution curve of  $u_x$  for the lighter gas is broader than that for the heavier gas.

(E) The root mean square of  $u_x$  is equal to  $(k_B T/m)^{1/2}$ .

(22) The rate of formation of  $\text{NH}_3$ , namely  $d[\text{NH}_3]/dt$ , in the reaction

$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$  was reported as  $1.2 \text{ mol L}^{-1} \text{ s}^{-1}$  (L is liter) under certain conditions. What is the



# 國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 7 頁

rate of consumption of H<sub>2</sub> gas, namely,  $\frac{d[H_2]}{dt} = ?$

(A) 1.2 mol L<sup>-1</sup> s<sup>-1</sup> (B) 1.4 mol L<sup>-1</sup> s<sup>-1</sup> (C) 1.8 mol L<sup>-1</sup> s<sup>-1</sup> (D) 2.4 mol L<sup>-1</sup> s<sup>-1</sup> (E) Not enough information.

(23) The gaseous isomerization reaction, CH<sub>3</sub>NC → CH<sub>3</sub>CN, displays first-order kinetics, namely rate =  $k[CH_3NC]$ . Measurements at 500 K show that the concentration of the reactant has declined to 75% of its initial value after 440 s. How much time will be required for the concentration of CH<sub>3</sub>NC to drop to 25% of its initial value?

(A) 640 s (B) 880 s (C) 1320 s (D) 1680 s (E) 2120 s.

(24) For a reversible reaction  $A \rightleftharpoons Y+Z$ , the rate constants  $k_1$  (forward) and  $k_{-1}$  (reverse) can be obtained by the T-jump relaxation method. Suppose  $x_e$  is the concentration of Y and Z at equilibrium. The relaxation time  $\tau =$

(A)  $\frac{1}{k_1 + k_{-1}}$  (B)  $\frac{1}{2k_{-1}x_e + k_1}$  (C)  $\frac{1}{2k_1x_e + k_{-1}}$  (D)  $\frac{1}{4k_1x_e + k_{-1}}$  (E)  $\frac{1}{4k_{-1}x_e + k_1}$

(25) In a consecutive first order reaction,  $A \rightarrow B \rightarrow C$  (where  $k_1$  and  $k_2$  are the respective rate constants), species B has transient existence. Therefore,

(A)  $k_1 \sim k_2$  (B)  $k_1 = 2k_2$  (C)  $k_1 \gg k_2$  (D)  $k_1 \ll k_2$  (E) None of these.

## 分析化學

單選題 (每題兩分，共 40 分)

(26) 50 mL of 0.1 M diprotic acid (H<sub>2</sub>A; pK<sub>a1</sub> = 10<sup>-4</sup>, pK<sub>a2</sub> = 10<sup>-8</sup>) reacted with 50 mL of 0.1 M NaOH. What is the pH of the resulting solution?

(A) 8.0 (B) 7.0 (C) 6.0 (D) 5.0 (E) 4.0

(27) What is the pH of 0.1 M NaCl solution? The activity coefficients of H<sub>3</sub>O<sup>+</sup> and OH<sup>-</sup> at ionic strength 0.1 M are 0.83 and 0.76.

(A) 7.00 (B) 7.01 (C) 7.02 (D) 6.99 (E) 6.98

(28) What are the ionic strength and pH of the buffer solution consisting of 0.1 M CH<sub>3</sub>COOH (pK<sub>a</sub> = 4.76) and 0.1 M CH<sub>3</sub>COONa?

(A) 0.1 M, pH 4.76 (B) 0.2 M, pH 5.76 (C) 0.3 M, pH 3.76 (D) 0.4 M, pH 2.76 (E) 0.6 M, pH 6.76

(29) The investigator developed the electrochemical biosensor for sensitive and selective detection of glucose. The obtained calibration curve for quantifying 1–100 mM glucose is  $y$  (signal) = 100 (slope, nA/mM) ×  $x$  (glucose concentration, mM) + 2. The standard deviation for measuring a blank solution (i.e., without glucose) is 1.5 nA. What is the limit of quantification of glucose measured by the electrochemical biosensors?

(A) 0.015 (B) 0.045 (C) 0.030 (D) 0.15 (E) 0.45.

(30) Which of the following conditions is (are) favorable for the formation of crystalline precipitation?

- (i) the use of a homogeneous precipitating agent,
- (ii) the increase in solution temperature,
- (iii) the dilution of a precipitating agent,
- (iv) the rapid injection of a precipitating agent,



國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 8 頁

(v) no stirring of the solution.

(A) i (B) i, ii (C) i, ii, iii (D) i, ii, iii, iv (E) i, ii, iii, iv, v

(31) What is the solubility of  $\text{Al}(\text{OH})_3$  ( $K_{sp} = 2.7 \times 10^{-35}$ ) in water?

(A)  $10^{-9}$  M (B)  $10^{-8}$  M (C)  $2.7 \times 10^{-14}$  M (D)  $2.7 \times 10^{-15}$  M (E)  $2.7 \times 10^{-35}$  M

(32) Which of the following statements associated with titration is (are) **CORRECT**?

(i) Titrations with NaOH are called argentometric titrations

(ii) In the Vohhard method,  $\text{Fe}^{3+}$  serves as the indicator

(iii)  $\text{Zn}^{2+}$  ions are titrated with EDTA in an ammonia/ammonium chloride buffer. The role of ammonia is to avoid the precipitation of  $\text{Zn}^{2+}$  ions.

(iv) The bromate titration method is commonly performed in a basic solution.

(A) i, ii, iii, iv (B) iii, iv (C) ii, iii, (D) ii, iii, iv (E) ii

(33) What is (are) the purpose of using high electrolyte solutions in electrochemical systems?

(i) To avoid the migration of ions close to the electrode surface by electrostatic attraction under applied voltage, which can lead to adsorption on the electrode surface.

(ii) To avoid too high resistance of the solution, which may cause IR drop.

(iii) Avoid redox reactions in the solution.

(iv) Effectively reduce the surface charge of the analyte, allowing the analyte to approach the electrode surface by diffusion and convection.

(A) i, ii (B) i, ii, iii (C) i, ii, iv (D) i, iii, iv (E) i, ii, iii, iv

(34) What approaches can be used to determine the molecular weight of an unknown protein?

(i) Matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS).

(ii) Size-exclusion chromatography coupled to UV absorption detector.

(iii) Capillary gel electrophoresis with UV absorption detector.

(iv) Gas chromatography-mass spectrometry (GC/MS).

(v) Thermogravimetric analyzer (TGA)

(A) i, ii, iii (B) i, ii, iii, iv (C) i, ii, iii, iv, v (D) i, ii, iv (E) i, ii, iii, v

(35) Which of the following statements associated with atomic spectroscopy is (are) **CORRECT**?

(i) Atomic absorption can be a thousand times more sensitive than atomic fluorescence.

(ii) For Beer's law to apply, the source must have a narrower linewidth than the atomic vapor.

(iii) In atomic absorption spectroscopy, nebulizers are used to introduce liquid samples.

(iv) An electrically heated graphite furnace is less sensitive than a flame burner and requires more samples.

(A) i, ii, iii, iv (B) i, ii, iii (C) ii, iii (D) iii, iv (E) i, iv

(36) Which of the following statements associated with X-ray fluorescence (XRF) is (are) **CORRECT**?

(i) XRF occurs when the material is excited by high-energy X-rays or gamma rays and emits characteristic secondary X-rays.

(ii) Elements and chemicals can be analyzed with XRF.

(iii) Planck's Law can be used to calculate the wavelength of the fluorescent radiation from an XRF-excited sample.

(iv) An electron from one of an atom's inner orbital shells is dislodged when an X-ray of sufficient energy strikes an atom in the sample.

(A) i (B) i, ii (C) i, ii, iii, (D) i, ii, iii, iv (E) i, iii, iv

(37) Which of the following statements associated with the ion-selective electrode is (are) **CORRECT**?

國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 9 頁

- (i) In place of aqueous NaCl in the salt bridge, the junction potential can be reduced to 0.1 mV.  
(ii) A typical pH combination electrode includes a working electrode, a glass membrane, a reference electrode, and a counter electrode.  
(iii) The boundary potential is the difference between the potential at the internal and external surface of the glass membrane.  
(iv) The sodium error of the pH meter occurs at low pH conditions.  
(A) iii (B) iii, iv (C) i, iii (D) ii (E) i, ii, iii, iv
- (38) Which of the following statements associated with redox titration is (are) **CORRECT**?  
(i) The titration of  $\text{Fe}^{3+}$  with  $\text{Ce}^{4+}$  can be recorded potentiometrically with Pt and calomel electrodes.  
(ii) The standard reduction potential of analytes can be determined by recording the potential change during redox titration.  
(iii) For permanganate titration,  $\text{KMnO}_4$  serves as an indicator and an oxidant in a basic solution.  
(iv) To titrate a reducing analyte with excess triiodide, thiosulfate is often used as a titrant for the remaining triiodide in a back titration.  
(A) i, ii, iii, iv (B) ii, iii, iv (C) iii, iv (D) i, iv (E) ii, iii
- (39) An electrode's overpotential is the voltage required to overcome its activation energy. Which of the following electrodes has the smallest overpotential?  
(A) Pt (B) Hg (C) Pb (D) Cu (E) Au
- (40) Coulometry is a chemical analysis based on counting the electrons used in a reaction. What is (are) the correct statement about coulometric titration?  
(i) The advantages of coulometry include precision, sensitivity, and generation of stable reagents in situ.  
(ii) Cyclohexane can be titrated with  $\text{Br}_2$  generated by electrolytic reduction of  $\text{Br}^-$ .  
(iii) Constant-current coulometry is performed using double Pt-wire electrodes.  
(iv) Controlled-potential coulometry in a three-electrode electrode cell is more selective than constant-current coulometry.  
(A) i, ii (B) iii, iv (C) i, ii, iii (D) i, iii, iv (E) i, ii, iii, iv
- (41) Which electrochemical analysis method is most sensitive to  $\text{Cd}^{2+}$  detection?  
(A) Linear scan voltammetry  
(B) Cyclic voltammetry  
(C) Square wave voltammetry  
(D) Anodic stripping analysis  
(E) Differential pulse voltammetry
- (42) Which of the following conditions can enhance the fluorescence of fluorescent molecules?  
(i) Enhancing fluorescent molecules' absorption at excitation wavelengths  
(ii) Utilizing an antireflective array to reduce fluorescent energy dissipation.  
(iii) Restricting the rotation of the fluorescent molecules in the viscous solution  
(iv) increasing the power of the excitation source  
(A) i, ii, iii, iv (B) i, ii, iii (C) ii, iii, iv (D) iii, iv (E) None of the above
- (43) Which of the following methods can enhance Raman signals?  
(i) The narrow linewidth of the laser is used in the Raman system.  
(ii) An analyte is absorbed on the roughened metal surface.  
(iii) Laser light is matched to the absorption maxima of the molecule.  
(iv) The charge-transfer process is conducted by adding semiconductor-based nanomaterials.  
(A) i, ii, iii, iv (B) i, ii, iii (C) ii, iii, iv (D) iii, iv (E) None of the above

# 國立中山大學 112 學年度碩士班暨碩士在職專班招生考試試題

科目名稱：物理化學及分析化學【化學系碩士班】

題號：422002

※本科目依簡章規定「可以」使用計算機（廠牌、功能不拘）（混合題）

共 10 頁第 10 頁

(44) Which element has an odd number of  $m/z$  for its parent peak in any compound?

(A) Nitrogen (B) Oxygen (C) Sulfur (D) Phosphorus (E) None of the above

(45) In your sample, the difference in mass between the two analytes is only 0.1  $m/z$ , which means 1140 and 1140.1. What is the minimum mass spectrometric resolution required?

(A) 1140 (B) 11400 (C) 114000 (D) 114 (E) None of the above

問答題 (每題 5 分，共 10 分)

(46) Fourier-transform infrared spectroscopy (FTIR) has several advantages over dispersive IR spectroscopy, including excellent spectral quality, high data collection speed, and high reproducibility. Explain why FT-IR has the above advantages

(47) Compare the external calibration curve method with the method based on standard additions and explain which is more accurate.