

**元智大學 102 學年度研究所 碩士班 招生試題卷**

系(所)別： **機械工程學系碩士班**      組別： **甲組**      科目： **工程數學**      用紙第 **1** 頁共 **2** 頁

●不可使用電子計算機

1. Using the method of variation of parameters to solve the differential equation. (16%)

$$y'' - \frac{4}{x}y' + \frac{4}{x^2}y = x^2 + 1, \quad x > 0$$

2. Using the method of Laplace Transformation to solve the initial value problem of  $y(t)$ . (17%)

$$y'' + 4y' + 3y = e^t \quad \text{with} \quad y(0) = 0, \quad \left. \frac{dy}{dt} \right|_{t=0} = 2$$

3. Given a curve  $C: \vec{r}(t) = \cosh t \vec{i} + \sinh t \vec{j}$ . (13%)

- (a) Find a tangent vector  $\vec{r}'(t)$  and the corresponding unit tangent vector  $\vec{u}(t)$ . (5%)
- (b) Find  $\vec{r}'(t)$  and  $\vec{u}(t)$  at the given point  $P: (\frac{5}{3}, \frac{4}{3}, 0)$ . (5%)
- (c) Find the tangent at  $P$ . (3%)

4. What kind of conic section (or pair of straight lines) is given by the quadratic form  $4x_1x_2 + 3x_2^2 = 1$ ? Transform it to principal axes. Express  $\mathbf{x}^T = [x_1 \quad x_2]$  in terms of the new coordinate vector  $\mathbf{y}^T = [y_1 \quad y_2]$ . (10%)

5. Using Green's theorem, evaluate the line integral  $\oint_C \vec{F}(\vec{r}) \cdot d\vec{r}$  counterclockwise around the boundary  $C$  of the region  $R$ , where  $\vec{F} = \left[ \frac{e^y}{x}, e^y \ln x + 2x \right]$ ,  $R: 1 + x^4 \leq y \leq 2$ . (10%)

**102023**

元智大學 102 學年度研究所 碩士班 招生試題卷

系(所)別： 機械工程學系碩士班


組別： 甲組

科目： 工程數學

用紙第 2 頁共 2 頁

●不可使用電子計算機

6. A two-dimensional rectangular plate is subjected to the boundary conditions shown as below. Derive an expression for the steady state temperature distributions  $T(x,y)$  with solving the heat conduction equation. (17 %)



The heat conduction equation is :

$$\frac{\partial^2 T(x,y)}{\partial x^2} + \frac{\partial^2 T(x,y)}{\partial y^2} = 0$$

Please find the solution in sin, cos, sinh, cosh series functions by the method of separation variables.

7. There is a partial differential equation of one dimensional wave equation to modeling vibrating string. The equation, boundary and initial conditions are following.

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}, u(0,t) = 0, u(l,t) = 0 \text{ for all time.}$$

The initial profile of the string is ,

$$f(x) = \begin{cases} \frac{2k}{l}x, & \text{when } 0 < x < l/2, \\ \frac{2k}{l}(l-x), & \text{when } l/2 < x < l \end{cases}$$

initial velocity is zero,  $g(x) = 0$ .

Please find the solution in sin, cos series functions by the method of separation variables. (17 %)

102039