淡江大學 100 學年度碩士班招生考試試題

系別:數學學系

科目:機

考試日期:2月28日(星期一) 第4節

大題,

頁

(10%) Suppose that A, B, and D are events such that A and B are independent,

$$P(A \cap B \cap D) = 0.04, P(D|A \cap B) = 0.25, \text{ and } P(B) = 4P(A).$$

Determine $P(A \cup B)$.

2. (10%) Let X and Y be independent random variables. Each has possible values 0, 1, and 2, with probabilities 0.2, 0.3, and 0.5, respectively.

a) (5%) Find the value of P(|X - Y| = 1).

b) (5%) Find P(X = x | Y = 1).

3. (20%) The sales of a convenience store on a randomly selected day are X thousand dollars, where X is a random variable with a distribution function of the following form:

$$F(t) = \begin{cases} 0 & t < 0, \\ t^2/2 & 0 \le t < 1, \\ k(4t - t^2) & 1 \le t < 2, \\ 1 & t \ge 2. \end{cases}$$

Suppose that this convenience store's total sales on any given day are less than 2000 dollars.

a) (5%) Find the value of k.

b) (15%) Let A and B be the events that tomorrow the store's total sales are between 500 and 1500 dollars, and over 1000 dollars, respectively. Are A and B independent events?

4. (15%) Suppose that three random variables X_1, X_2 , and X_3 have a continuous joint distribution with the following joint probability density function

$$f(x_1, x_2, x_3) = \begin{cases} ce^{-(x_1 + 2x_2 + 3x_3)} & \text{for } x_i > 0 \ (i = 1, 2, 3), \\ 0 & \text{otherwise.} \end{cases}$$

Determine

is

a) (5%) the value of the constant c;

b) (10%) $P(X_1 < 1 | X_2 = 2, X_3 = 1).$

(15%) Suppose that X is a random variable for which the moment generating function is

$$\phi(t) = \frac{e^t}{5} + \frac{2e^{4t}}{5} + \frac{2e^{8t}}{5}$$
 for $-\infty < t < \infty$.

a) (5%) What is the probability mass function of X?

b) (10%) Compute Var(X).

(15%) Suppose that the joint probability density function of two random variables X and Y

$$f(x,y) = \frac{1}{2\pi} e^{-(x^2 + y^2)/2}$$
 for $-\infty < x < \infty$ and $-\infty < y < \infty$.

Find $P(-\sqrt{2} < X + Y < 2\sqrt{2})$ in terms of $\Phi(z)$, where $\Phi(z) = \int_{-\infty}^{z} \frac{1}{\sqrt{2\pi}} e^{-\frac{w^2}{2}} dw$.

7. (15%)Let X_1, X_2, \ldots, X_n be independent random variables, each having the uniform distribution on (0, 1). Let $Y = -2\ln(X_1X_2\cdots X_n)$. What is the distribution of Y?